



STUDENT LEARNING OUTCOMES [C-11-B-30 to C-11-B-41]

- Explain the lack of reactivity of nitrogen due to its triple bond strength and lack of polarity. **(Understanding)**
- Describe the basicity of ammonia using the Bronsted-Lowery theory. **(Understanding)**
- Identify the structure of the ammonium ion and explain how it is formed by an acid-base reaction. **(Understanding)**
- Describe how ammonia can be displaced from ammonium salts through an acid-base reaction. **(Understanding)**
- Describe natural and man-made occurrences of oxides of nitrogen and their catalytic removal from exhaust gases of internal combustion gases. **(Understanding)**
- Explain the role of NO and NO₂ in the formation of photochemical smog, specifically in the reaction with unburned hydrocarbons to form peroxyacetyl nitrate (PAN). **(Understanding)**
- Differentiate between nitrification and de-nitrification. **(Knowledge)**
- Explain the lack of reactivity of sulfur, with reference to its bonding and stability of its compounds. **(Understanding)**
- Describe the different oxidation states of sulfur and their relative stability. **(Understanding)**
- Describe the properties and uses of sulfuric acid, including its production and industrial applications. **(Understanding)**
- Describe the chemical reactions and processes involving sulfur, such as combustion and oxidation. **(Understanding)**
- Explain the uses of sulfur compounds in industry and everyday life, such as in fertilizers, gunpowder, and rubber, and in synthetic organic chemistry, including the synthesis of dyes, drugs, and fragrances. **(Understanding)**

NITROGEN

Nitrogen belongs to group 15 of the periodic table. In industrial processes, nitrogen is typically obtained by cooling air until it becomes a liquid. Liquid nitrogen is commonly used for rapid cooling purposes. In laboratory settings, nitrogen can be generated by slowly heating a solution of ammonium nitrite.




Table 12.1 Physical properties of nitrogen

Atomic number	7	Ionic radius (3-)	171 pm
Relative atomic mass	14.007 Da	1 st Ionization energy	1402 kJ/mol
Physical appearance	Colourless Gas	Electronegativity	3.0
Electronic configuration	[He] 2s ² 2p ³	Electron affinity	-8.0 kJ/mol
Melting point	-210 °C	Density	0.001145 g/cm ³
Boiling point	-195.8 °C	Principal oxidation states	3+ and 5+
Covalent radius	74 pm		

12.1 REACTIVITY OF NITROGEN (N₂)

Nitrogen is a significant component of the air, known for its low reactivity due to its small size, symmetrical electronic cloud, and nonpolar triple bond. With an electronic configuration of 1s² 2s² 2p³, nitrogen requires three electrons to complete its octet, forming a triple bond by sharing three electrons with another nitrogen atom as shown in **Figure 12.1**. This bond has a bond enthalpy of +944 kJmol⁻¹. High energy is required to break this bond to form new bonds, making N₂ very unreactive. The second reason for its lack of reactivity is the non-polarity of its bond. Both the atoms are the same having zero electronegativity difference. This causes equal sharing of the three bonded electrons between the two atoms making the bond nonpolar.

Figure 12.1 Bonding in N₂ molecule

Interesting Information!

The high concentration of nitrogen in the air serves to dilute oxygen, preventing every spark in our atmosphere from igniting a massive fire. In the case of large shipments of hydrocarbons or edible oils, it is crucial to utilize blankets of Nitrogen or any other inert gas on ships to safeguard them from oxygen and moisture. It is also used in laboratory to carry out the reactions which require inert atmosphere.

12.2 AMMONIA (NH₃)

Ammonia (NH₃) is an important industrial compound of nitrogen, which is mainly used as a fertilizer. It is prepared industrially by Haber-Bosch process.




12.2.1 Basicity of Ammonia

Ammonia behaves as a Lowry-Bronsted base by accepting a proton (H⁺) from an acid to form ammonium:



It dissolves in water to form ammonium hydroxide (NH_4OH) and equilibrium is established between ammonia molecules and ammonium ions in the solution.



Ammonia solution is a weak base due to the low basicity constant (K_b) and the equilibrium position being towards the far left side.

12.2.2 Structure of Ammonium (NH_4^{1+})

Ammonia molecule has pyramidal shape due to lone pair of nitrogen. But when nitrogen atom in ammonia utilizes this lone pair of electrons to form ammonium, this ion adopts a tetrahedral shape in which all the bonds are of equal length and strength, as depicted in **Figure 12.2**.





Figure 12.2 Pyramidal and tetrahedral shapes of ammonia and ammonium

12.2.3 Synthesis of Ammonia from Ammonium salts

In the laboratory, ammonia gas can be synthesized by heating an ammonium salt such as ammonium chloride (NH_4Cl) with a base like calcium hydroxide ($\text{Ca}(\text{OH})_2$) as shown in **Figure 12.3**.



In this acid-base reaction, NH_4^+ acts as an acid by donating H^+ ions, while OH^- acts as a base by accepting H^+ ions. This reaction displaces ammonia gas from the ammonium salt and produces salt and water. It is commonly used to identify ammonium in salt analysis. If a gas with a pungent smell is released and turns moistened red litmus paper blue, it indicates the presence of ammonium in the compound.




Figure 12.3 Synthesis of ammonia gas

Quick Check 12.1

- Why N_2 gas is used in food packaging?
- Both CO and N_2 have triple bonds in their molecules. Why do you think CO is more reactive than N_2 ?
- Ammonium salts such as $(\text{NH}_4)_2\text{SO}_4$ or $(\text{NH}_4)_2\text{NO}_3$ are commonly used as fertilizers. Why a farmer wouldn't treat a field with an ammonium fertilizer at the same time as using lime? What would be the chemical reactions?
- If magnesium ribbon is ignited and placed in a jar containing N_2O , it continues to burn brightly. Explain this observation with reason.

12.3 OXIDES OF NITROGEN

Oxides of nitrogen are NO , N_2O , NO_2 , N_2O_4 and N_2O_5 in which oxidation states range from I^+ to V^+ . N_2O_4 and N_2O_5 decay quickly to other oxides. NO and NO_2 are collectively called as NO_x . The structures, properties and uses of these oxides are given in **Table 12.2**.

Table 12.2 Properties of some common oxides of nitrogen

Name and formula of oxide	Structure	Formal oxidation state	Properties	Uses
Nitrous oxide/ Nitrogen oxide (Laughing gas) N_2O	$\text{:N}\equiv\text{N}-\ddot{\text{O}}\text{:}$	1^+	Colourless gas, water-soluble, neutral, sweet smelling, helps in combustion	Dental anaesthetic, propellant for whipped ice cream, synthesis of NaN_3
Nitric oxide, Nitrogen dioxide NO , N_2O_2	$\text{:N}\equiv\ddot{\text{O}}$ $\text{:}\ddot{\text{O}}=\ddot{\text{N}}-\ddot{\text{N}}=\ddot{\text{O}}\text{:}$	2^+	Colourless gas, slightly water-soluble, neutral, NO is paramagnetic while N_2O_2 is diamagnetic, oxidising as well as reducing in nature	Biochemical messenger (Lowers blood pressure and role in other body functions), synthesis of nitrosyl carbonyls
Nitrogen dioxide/Nitrogen peroxide, Nitrogen tetraoxide NO_2 , N_2O_4 $2\text{NO}_2 \xrightleftharpoons[\text{Heat}]{\text{Cool}} \text{N}_2\text{O}_4$ Brown Colourless	$\text{:}\ddot{\text{O}}-\ddot{\text{N}}=\ddot{\text{O}}\text{:}$ $\text{:}\ddot{\text{O}}\text{:}\text{N}(\text{:}\ddot{\text{O}}\text{:})_2\text{N}(\text{:}\ddot{\text{O}}\text{:})_2\text{:}$	4^+	NO_2 is a reddish-brown gas, paramagnetic, and reacts with water to form HNO_3 and HNO_2 , N_2O_4 is colourless liquid or solid	Rocket propellant, HNO_3 formation by Ostwald process, explosives



Did you know?

Another recommended name for ammonia by the International Union of Pure and Applied Chemistry (IUPAC) is *Azane* and for ammonium is *Azanium*. These names have been derived from *Azote*, a Greek name for nitrogen, meaning “no life”. These names are used in naming derivatives of ammonia and ammonium e.g. sodium azide (NaN_3).

12.4 SOURCES OF OXIDES OF NITROGEN

Here are the main categories of NO_x sources.

12.4.1 Natural Sources

Natural sources include lightning, volcanoes, biological decay, forest fires, soil microorganisms, oceans, etc. NO is produced when N_2 and O_2 in the air react during lightning. It is produced by microorganism using air N_2 .

12.4.2 Anthropogenic (Man-made) Sources





The main anthropogenic sources of NO_x are the combustion of fossil fuels in vehicles and power plants. Other sources include chemical plants, biomass burning, welding, etc.

12.5 ROLE OF NO & NO_2 IN SMOG & PAN FORMATION

NO_x is responsible for numerous harmful effects on living organisms.

12.5.1 Photochemical Smog

Photochemical smog (Los Angeles smog) forms in the atmosphere from NO_x and volatile organic compounds (VOCs) in the sunlight. It is oxidising in nature. Photochemical oxidants, such as NO_2 , ozone, and peroxyacyl nitrates (PANs) can react and oxidise specific compounds in the atmosphere. Photochemical smog is becoming more common than classical smog (London smog) due to increasing NO_x emissions. The formation of photochemical smog involves the following chemical reactions.



Did you know?

Lahore smog consists of volatile organic compounds (VOCs), NO_x , ground level ozone (O_3), particulate matter $\text{PM}_{2.5}$, CO and SO_2 .

12.5.2 Formation of Peroxyacyl Nitrates (PANs)

NO_x take part in a series of reactions leading to the formation of ozone (O_3), aldehydes, peroxyacyl nitrates (PANs) and peroxybenzoyl nitrate (PBN). PAN is one of the members of peroxyacyl nitrates as shown in **Figure 12.4**.




Figure 12.4 General structure of PANs

The R group in peroxyacetyl nitrate is CH_3 , but other hydrocarbon chains may also be present.

The formation of PAN is illustrated in **Figure 12.5**. The main component of oxidizing smog is ozone, which oxidizes hydrocarbon to produce aldehyde. The aldehyde then reacts with hydroxyl radical to produce acyl radical. The acyl radical reacts with oxygen to produce peroxyacyl radical, which finally reacts with nitrogen peroxide to form peroxyacyl nitrate.




Figure 12.5 Mechanism of PAN formation

Quick Check 12.2

- Draw the structures of the following oxides of nitrogen. Also, briefly explain their bonding.
 - N_2O
 - NO
 - NO_2
- What does PAN stand for? Give its general formula.
- Write down the formulas of the compounds responsible for the formation of PAN.
- If magnesium ribbon is ignited and placed in a jar containing N_2O , it continues to burn brightly, how does the product form in this reaction confirm the structure of N_2O ?


12.6 CATALYTIC CONVERTER

A catalytic converter is a ceramic or metallic monolith with a honeycomb-like structure, as shown in **Figure 12.6**. Its inner channels have a layer of alumina to provide a high surface area. Noble expensive metals such as Pt (Platinum), Pd (Palladium), and Rh (Rhodium) are

dispersed on the alumina.

These metals catalyze three redox reactions to remove the half harmful exhaust gases. The three-way converter converts harmful CO, NO, and hydrocarbons into CO₂, N₂, and water. These precious metals can also be recycled.

Reduction



Oxidation






Figure 12.6 Three-way catalytic converter

12.7 NITRIFICATION AND DENITRIFICATION

Nitrification and denitrification are the two phases of the nitrogen cycle. Nitrification involves the conversion of ammonium (NH₄¹⁺) to nitrite (NO₂¹⁻) and nitrate (NO₃¹⁻), while denitrification involves the conversion of nitrate (NO₃¹⁻) to Nitrogen (N₂). These two processes are also involved in the wastewater treatment to remove nitrogen. Some differences between nitrification and denitrification are given in Table 12.2.

Table 12.3 Differences between nitrification and denitrification

Nitrification	Denitrification
Ammonia NH ₃ /NH ₄ ¹⁺ is converted into nitrite (NO ₂ ¹⁻) and nitrate (NO ₃ ¹⁻).	Nitrite (NO ₂ ¹⁻) and nitrate (NO ₃ ¹⁻) are converted back to N ₂ that is released into the atmosphere.
Nitrifying bacteria aerobic conditions, pH 6.5 – 8.0, optimum temperature 20 °C–30 °C.	Denitrifying bacteria anaerobic conditions, pH 7.0–9.0, optimum temperature 26 °C – 38 °C.
Plants absorb these nitrites and nitrates for their nutrition as they cannot assimilate nitrogen directly from the atmosphere.	It is important in wastewater treatment and useful for aquatic life, to oxidize NH ₄ ⁺ with NO ₂ ⁻ to form N ₂ gas.
Oxidation of nitrogen $\text{NH}_4^+ \rightarrow \text{NO}_2^- \rightarrow \text{NO}_3^-$ $2\text{NH}_{4(\text{aq})}^{1+} + 3\text{O}_{2(\text{aq})} \rightarrow 2\text{NO}_{2(\text{aq})}^{1-} + 4\text{H}_{(\text{aq})}^+ + 2\text{H}_2\text{O}_{(\text{l})}$ $2\text{NO}_{2(\text{aq})}^{1-} + \text{O}_{2(\text{aq})} \rightarrow 2\text{NO}_{3(\text{aq})}^{1-}$	Reduction of nitrogen $\text{NO}_3^- \rightarrow \text{NO}_2^- \rightarrow \text{NO} \rightarrow \text{N}_2\text{O} \rightarrow \text{N}_2$ $2\text{NO}_{3(\text{aq})}^{1-} + 10\text{e}^- + 12\text{H}_{(\text{aq})}^+ \rightarrow \text{N}_{2(\text{g})} + 6\text{H}_2\text{O}_{(\text{l})}$

Quick Check 12.3

- Write down the reduction and oxidation reactions that occur in the catalytic converter in the vehicle exhausts.
- What is the basic principle of catalytic converter? Describe the role of catalyst in the catalytic converter.
- Do hybrid and electric cars have catalytic converters? Explain why or why not.

12.8 SULFUR


Sulfur is a member of group 16 which is also called the *Chalcogen* family. Some physical properties of sulfur are listed in **Table 12.3**.

Table 12.4 Physical properties of Sulphur

Atomic number	16	Ionic radius	184 pm
Relative atomic mass	32.06 a.m.u.	1st Ionization energy	1000 kJ/mol
Physical appearance	Solid Yellow	Electronegativity	2.5
Electronic configuration	[Ne] 3s ² 3p ⁴	Electron affinity	-200 kJ/mol
Melting point	113 °C (honey-yellow)	Density	2.07 g/cm ³
Boiling point	445 °C (dark brown)	Common oxidation states	4+ and 6+
Covalent radius	104 pm	Common Bonding	Covalent bond

12.8.1 Reactivity of Sulfur

Sulfur usually forms single bonds with other sulfur atoms instead of double bonds due to poor overlapping of the orbitals. As a result, it forms larger molecules and structures through a process called catenation. S₈ is a crown-like molecule, as shown in **Figure 12.7**.

**Figure 12.7** Cyclo-octasulfur, (S₈) crown molecule

12.8.2 Oxidation States of Sulfur

Sulfur exhibits oxidation states of -2, 0, +2, +4, and +6. The electronic configuration of sulfur is depicted in **Figure 12.8**, where the oxidation state is determined by the number of unpaired electrons. Under standard conditions, sulfur and oxygen react to produce sulfur

dioxide (SO_2) in which sulphur has an oxidation state of +4. However, to form sulfur trioxide (SO_3) with an oxidation state of +6, high energy is required.







a) Oxygen atom in ground state ($2s^2 2p^4$)	$2s$ 	$2p$ 	There are no d-orbitals in 2 nd shell therefore excitation is not possible.	
b) Sulfur atom in ground state ($3s^2 3p^4$)	$3s$ 	$3p$ 	$3d$ 	2+ oxidation state due to two unpaired electrons
c) Sulfur atom in 1st excited state ($3s^2 3p^3 3d^1$)				4+ oxidation state due to four unpaired electrons
d) Sulphur atom in second excited state ($3s^1 3p^3 3d^2$)				6+ oxidation state due to six unpaired electrons

Figure 12.8 Electronic configurations of sulfur for attaining different oxidation states

12.9 STABILITY OF OXIDATION STATES OF SULPHUR

The Stability of oxidation states of sulphur is explained by Frost diagram in which an element X is a plot of nE° vs oxidation state (n) for the redox couple $X(n)/X(0)$ as shown in **Figure 12.9**. It tells about the relative stabilities of different oxidation states of an element at two extreme pH values (pH=0 and pH=14) i.e. acidic and basic conditions. It can also be regarded as a plot between standard reaction Gibbs free energy (ΔG°) versus oxidation state (n) because oxidation state(n) is directly proportional to ΔG° as shown below.

$$-\Delta G^\circ = nFE^\circ$$

$$\text{or } \frac{-\Delta G^\circ}{F} = nE^\circ$$

Here E° is the standard reduction potential, F is the Faraday constant and n is the number of transferred electrons in a redox half reaction.

The more stable oxidation state has a lower nE° value or more negative Gibbs free energy. Therefore, the more stable oxidation state is at the lowest position in the diagram. Further, a couple with a more positive slope is a good oxidising agent, while a couple with a more negative slope is a good reducing agent.

In the diagram for sulfur, +6 (SO_4^{2-}) is the most thermodynamically stable oxidation state, completely non-oxidising under extreme basic conditions, while under extreme acidic conditions, it is shown to be very unstable.




Figure 12.9 Frost diagram for oxidation states of sulfur

It can be seen that in basic environments, the oxidized sulfur states are all more stable than the elemental form but under acidic conditions all the oxidized sulfur states are less stable than the elemental form.


Quick Check 12.4

- Determine the oxidation state of S in the following species.
i. SO_3 ii. H_2SO_4 iii. SO_4^{2-} iv. $\text{S}_2\text{O}_3^{2-}$
- Which oxidation states of sulphur are the most common? Explain your answer with reason.
- Explain the involvement of d orbital in variable oxidation states of S in its compounds.

12.10 REACTIONS OF SULFUR

Sulfur can combine with many elements to form a wide variety of inorganic and organic compounds. It is unreactive to water under normal conditions, dilute non-oxidising acids, and noble gases. Its ability to catenate allows it to form ring structures and linear chains. Here are some important reactions of sulfur:


- Sulfur burns in the air to form SO_2 with a blue colour flame. The other main sulphur oxide is SO_3 which requires higher temperature and a catalyst for its formation.



Sulfur can be oxidised by nitric acid to produce NO_2 and H_2SO_4 .




- When sulfur reacts with elements that have lower electronegativity, it acts as an oxidizing agent and forms their sulfides. It tarnishes Ag, Cu, and Zn by forming a coating of metal sulfide. Sulfur does not react with Au and Pt.



- Sulfur converts cyanide into thiocyanate which is also known as pseudohalide. It is used to analyse Fe^{3+} .



- Sulfur reacts directly with F_2 to form SF_4 and SF_6 . Sulfur hexafluoride (SF_6) is a gas and is very unreactive. It is used as an insulator gas in electric devices. Sulfur readily reacts with Cl_2 to form S_2Cl_2 (yellow liquid) which further reacts with Cl_2 to form SCl_2 (red liquid).



12.11 USES OF SULFUR AND ITS COMPOUNDS

12.11.1 Vulcanisation

Sulphur is used as a cross linker for the rubber molecular chains. This is called vulcanisation and it improves the strength of rubber as shown in **Figure 12.10**.

12.11.2 Fertilizer

Sulfur is an essential nutrient for plant growth. When soils become depleted in sulfate, sulfur can be restored in soil by applying sulfur containing N/P fertilizers, or sulfur-coated fertilizers such as sulfur-coated urea. Soil components and microbes convert elemental sulfur into soluble forms for the use of plants. Gypsum ($\text{CaSO}_4 \cdot 5\text{H}_2\text{O}$) is also used as a fertilizer.





Figure 12.10 S-S cross-linkages between polymer chains

12.11.3 Gun powder

Gun powder is a coarse blend of 75% potassium nitrate (KNO_3), 15% wood charcoal, and 10% sulfur. Charcoal carbon is the main fuel, nitrate is the oxidiser and sulfur is the additional fuel that burns the powder faster. The following reaction in burning takes place:



Quick Check 12.5

- What is the function of SO_2 and sulfite (SO_3^{2-}) salt in preserving the food?
- Give the reaction of S with HCl acid and NaOH.

12.12 ROLE OF SULFUR IN ORGANIC SYNTHESIS OF DRUGS, DYES AND FRAGRANCES

Carbon-Sulfur bonds are prevalent in a wide range of compounds with biological, pharmaceutical, and material properties. These bonds form a large number of organic compounds containing a variety of functional groups such as thiols or mercaptans, thioethers, sulfoxides, sulfones, etc.

12.12.1 Drugs

Sulfa drugs are the antibacterial sulfonamides such as penicillins and cephalosporins contain sulfur. The common drug omeprazole, used in GERD (Gastroesophageal reflux disease) contains sulfoxide group.

12.12.2 Dyes

Sulfur dyes are synthesised by the process of thionation or sulfurization of organic compounds that contain nitro or amino groups. These compounds contain sulfur linkages. They generally give black, brown, khaki, blue, and green colours.

Some examples are sulfur black 1, sulfur blue and sulfur brilliant green.

12.12.3 Odorants/Fragrances

Mercaptans are used to give odour to natural gas. Some thiols have pleasant odours on high dilution, for example, thioterpineol is the key ingredient in the aroma of grapefruit. cis-glabanum oxathiane is a fragrant compound. It is used in fine fragrances, soaps, shampoos and shower gels. Many naturally occurring odorants are produced synthetically and also applied as flavouring agents.

12.13 SULPHURIC ACID (H_2SO_4)

The major portion of sulfur, around 85% is used for the production of sulphuric acid (H_2SO_4). It has tetrahedral structure with two S–O and two S=O bonds as shown in **Figure 12.11**.




Figure 12.11 Structure of H_2SO_4

12.13.1 THE CONTACT PROCESS

Sulphuric acid is produced by the contact process. A flow sheet diagram of the contact process is shown in **Figure 12.12**.






Figure 12.12 Contact process for the industrial production of sulphuric acid

The Contact process can be divided into the following stages.


Sulfur/pyrite Burners

The process starts with the combustion of molten sulphur or by heating pyrites such as iron pyrite (FeS) in excess of air to produce sulfur dioxide SO₂.



Purification Unit

If pyrite ore is used as a sulfur source, the SO₂ gas formed may contain contaminants like dust particles, vapours, and arsenic oxide. These contaminants affect the efficiency of the catalyst. Hence, the gas needs to pass through the purification unit. In an arsenic purifier, gelatinous ferric hydroxide Fe(OH)₃ present in horizontal shelves, absorbs arsenic oxide As₂O₃.




Contact Tower and Heat Exchangers

Purified SO₂ and air, preheated at 420 °C– 450 °C, are fed to the first converter stage of the contact tower at 1–2 atm pressure. Here, these gases come in contact with vanadium pentoxide (V₂O₅) catalyst.




The catalyst works in two steps:

Oxidation of SO₂ into SO₃ by V⁵⁺:



Oxidation of V⁴⁺ back into V⁵⁺ by oxygen (catalyst regeneration):




Quick Check 12.6

- Why is high temperature and catalyst needed to form SO₃?
- Write down dehydration reactions of conc. H₂SO₄ with starch and oxalic acid
- How does the catalyst V₂O₅ function in the conversion of SO₂ to SO₃?



Absorption Tower

Sulfur trioxide is cooled and can be converted to sulphuric acid by reacting with water.



Mixing SO₃ with water is not feasible because the reaction is extremely exothermic and acidic vapour or mist is produced rather than a liquid solution. Mainly, sulfur trioxide is dissolved in recirculating hot 98.5% sulphuric acid. The term fuming sulphuric acid or oleum is used for the mixtures of sulfur trioxide with 100 percent sulphuric acid.

Oleum undergoes a reaction with water to make a highly concentrated solution of H_2SO_4 whose concentration can be adjusted.



12.13.2 Physical Properties

Sulphuric acid is soluble in water and hygroscopic in nature. It readily absorbs water vapour from the air. Anhydrous H_2SO_4 is a very polar liquid. It is highly corrosive to various materials. On contacting the skin, it causes chemical burns. Some physical properties of sulphuric acid are given in **Table 12.4**.


Table 12.5 Physical properties of Sulphuric acid

Molar mass	98.08 g/mol
Physical appearance	colourless viscous liquid
Odour	odourless
Melting point	10 °C
Boiling point	290 °C
Specific gravity (15 °C)	1.83 g/cm ³
Viscosity	25.24 centipoise
Vapour pressure (25 °C)	0.001 torr
Sulphuric acid is a highly corrosive substance. It can badly burn cloths, plastic, rubber and injured the human skin, eyes, etc., Handle it very carefully.	




12.13.3 CHEMICAL PROPERTIES

i. It self-ionizes or undergoes autoprotolysis as follows.




The equilibrium constant value is greater than that of water which makes it to be used as a non aqueous protic solvent.

Sulphuric acid is a strong acid as shown by its pK_{a1} value:




But hydrogensulfate (HSO_4^{1-}) is a far weaker acid due to a positive pK_{a2} value:




ii. Concentrated sulphuric acid is a powerful dehydrating agent that removes water from many substances such as sucrose, starch, wood, and paper to produce carbon, steam, and heat.

**Interesting Information!**


A common laboratory demonstration is the dehydration of table sugar, where a black porous carbon mass called carbon snake protrudes out of the apparatus.




It also dehydrates ethyl alcohol to ethene or ethoxyethane depending upon the reaction conditions.



- iii. Hydrochloric acid (HCl) gas, is formed when sulphuric acid reacts with sodium chloride



- iv. Reactions of sulphuric acid with metals depend upon the metal, concentration of the acid, and temperature. Metals that are above hydrogen in electrochemical series such as Fe, Al, Zn, Mn, Ni, and Mg react directly with dilute sulphuric acid to produce hydrogen gas and metal sulfates. But with cold conc. H_2SO_4 , they liberate SO_2 and formsulfates.



- v. Metals like Cu, Ag, and Hg react with hot conc. H_2SO_4 to form metal sulfates. Sulphuric acid is not regarded as a typical oxidising agent due to the stability of SO_4^{2-} anion. This anion is weakly oxidising. However hot concentrated sulphuric acid is a moderately strong oxidising agent due to high temperature, high concentration of protons (H^+), and formation of nascent oxygen. Hot concentrated sulphuric acid oxidises Cu, as given below:

**Quick Check 12.7**

- Write down dehydration reactions of conc. H_2SO_4 with starch and oxalic acid.
- How does conc. H_2SO_4 react with NaCl? What is the importance of this reaction?
- What is the difference between the oxidizing power of cold and hot sulphuric acid?

12.13.4 Uses and Industrial Applications

Sulphuric acid is considered a king of chemicals and its consumption is an indicator of the industrial progress of a country.

- A major portion of the acid is used in making fertilizers, normally 77.67 % is used to digest the phosphate rock containing calcium phosphate ($\text{Ca}_3(\text{PO}_4)_2$).
- It is used in the extraction of metals from ores such as Cu, Ni, steel etc.
- It is utilized as a catalyst in oil and coal refining, polymers, synthetic rubber, and plastic industries.

4. It is used in the pulp and paper industry and involved in the production of pesticides, insecticides, herbicides, varnishes, dyes, pharmaceuticals, soaps and detergents.
5. It is used for nitration in making explosives such as trinitrotoluene (TNT), nitroglycerine, picric acid, nitrocellulose, etc.
6. It is involved in the food industry for making sugar, starch, and corn syrup.
7. It is used in the paint industry for making titanium dioxide (TiO_2) pigment.
8. It is used to dry gases in industrial processes.
9. 35.67% acid is used in lead storage batteries.
10. It is used as a laboratory reagent.

Even though it is used in various industries, it is rarely contained in the final product.



Did You Know?

Fluorapatite ($\text{Ca}_5(\text{PO}_4)_3\text{F}$) is used to make single superphosphate $\text{Ca}(\text{H}_2\text{PO}_4)_2$ and phosphoric acid (H_3PO_4). H_3PO_4 is further used to make double and triple superphosphates and ammonium phosphate. Sulphuric acid also reacts with ammonia to make ammonium sulfate fertilizer.

EXERCISE

MULTIPLE CHOICE QUESTIONS

Q.1 Four choices are given for each question. Select the correct choice.

I. Despite being the most abundant gas in the Earth's atmosphere, nitrogen does not readily participate in combustion reactions because:

- a) It is denser than oxygen.
- b) It has a high specific heat capacity.
- c) Breaking the $\text{N}\equiv\text{N}$ bond requires a large amount of energy.
- d) It is a noble gas.

II. A student heats a solid ammonium salt with a solution of a strong alkali. The gas produced turns damp red litmus paper blue and has a characteristic pungent smell. The gas is:

- | | |
|------------------------------|-------------------------------------|
| a) Hydrogen (H_2) | b) Carbon dioxide (CO_2) |
| c) Ammonia (NH_3) | d) Sulfur dioxide (SO_2) |

III. The shape of ammonium is

- | | |
|----------------|----------------------|
| a) pyramidal | b) triangular planar |
| c) tetrahedral | d) linear |

IV. In a catalytic converter, the conversion of nitrogen oxides (NO_x) into nitrogen gas (N_2) and oxygen gas (O_2) is a process of:

- a) Oxidation.
- b) Reduction.
- c) Combustion.
- d) Neutralization.

V. PAN formation starts when _____ reacts with the hydrocarbon.

- a) NO
- b) NO_2
- c) O_3
- d) $\text{HO}^\bullet$

VI. Nitrification is the process by which:

- a) Atmospheric nitrogen is converted into ammonia.
- b) Nitrate is converted into nitrogen gas.
- c) Ammonia is converted into nitrite and then nitrate.
- d) Organic nitrogen is converted into ammonia.

VII. The most stable species in an acidic environment is

- a) SO_4^{2-}
- b) SO_2
- c) H_2S
- d) S

VIII. Which gas is used in separating hard water from normal water?

- a) SO_2
- b) H_2S
- c) NH_3
- d) NO_2

IX. The oxidation state of sulfur in H_2SO_4 is

- a) +1
- b) +2
- c) +4
- d) +6

X. In a Frost diagram, a species that lies above the line connecting its neighbors is:

- a) Thermodynamically stable.
- b) A strong oxidizing agent.
- c) Prone to disproportionation.
- d) A poor reducing agent.

XI. Sulfur dioxide (SO_2) produced from the combustion of sulfur can be further oxidized to sulfur trioxide (SO_3) under specific conditions, such as in the presence of a:

- a) Catalyst (e.g., vanadium(V) oxide) and high temperature.
- b) Catalyst (e.g., iron) and low temperature.
- c) Strong reducing agent and high pressure.
- d) Dilute acid and room temperature.

XII. Sulfur trioxide (SO_3) is not directly dissolved in water to produce sulfuric acid in the Contact Process because this reaction is:

- a) Too slow.
- b) Reversible and would result in a low yield.
- c) Highly exothermic and produces a mist of sulfuric acid.
- d) Requires very high pressures.

SHORT ANSWER QUESTIONS

Q.2 Attempt the following short-answer questions:

- a. List two reasons for the inertness of N_2 .
- b. How is nitrogen isolated from air?
- c. Why ammonia (NH_3) is a weak base?
- d. How does NO_2 catalyze the formation of SO_3 ?
- e. Write down the reactions of photochemical smog formation.
- f. What is the construction and function of a catalytic converter?
- g. Why sulfur is quite unreactive at room temperature?
- h. Which are the most stable oxidation states of sulphur in water at $\text{pH}=0$ and $\text{pH}=14$?
- i. How does sulfur react with halogens?
- j. Draw the structures of cyclo-octasulfur (S_8) and sulfuric acid.
- k. What is the role of sulfur in the vulcanization of rubber?
- l. What is the composition and the chemical reaction of gunpowder combustion?
- m. What is the importance of disulfide bridges?
- n. Write down self-ionization equation of sulfuric acid and its ionization in water.
- o. Give two examples where sulfuric acid acts as a dehydrating agent.
- p. How does V_2O_5 catalyze the formation of SO_3 ?
- q. What is purpose of formation of oleum?

DESCRIPTIVE QUESTIONS

Q.3 Explain the preparation and basicity of ammonia.

Q.4 How oxides of nitrogen (NO_x) cause the formation of photochemical smog and PAN? Give its mechanism.

Q.5 Frost diagram explains the stabilities of different oxidation states of sulfur. Which oxidation states of sulphur are the most stable at acidic and basic conditions?

Q.6 Discuss sulfuric acid as an oxidizing agent and a dehydrating agent with three reactions for each.